


R

تداسپیت

متوپرولول سوکسینات (پیوسته رهش)



پیش از شصت سال تلاش در تعالی فرهنگ سلامت کشور



موثر در درمان

- * آنژین پکتوریس
- * نارسایی قلبی
- * فشار خون بالا
- * فشار خون بالا در کودکان بالای ۶ سال


مصارف Off-label:

- * پیشگیری از انفارکتوس مجدد میوکارد
- * کنترل سرعت ضربان بطنی در بیماران مبتلا به فیبریلاسیون دهلیزی
- * درمان نگهدارنده تاکی کاردی فوق بطنی
- * درمان مشکلات تیروئیدی (تیروتوکسیکوزیس)
- * پیشگیری از آریتمی بطنی
- * پیشگیری از سردردهای میگرنی



Tedacept[®]

Metoprolol Succinate (Extended-Release)



Pharmacologic Category:

Antianginal Agent; Antihypertensive; Beta-Blocker, Beta-1 Selective

Mechanism of Action:

Selective inhibitor of beta1 - adrenergic receptors; competitively blocks beta1-receptors, with little or no effect on beta2-receptors at oral doses <100 mg (in adults); does not exhibit any membrane stabilizing or intrinsic sympathomimetic activity.

Indications:

Angina Pectoris : Initial: 95 mg once daily; may increase dose at weekly intervals to desired effect; maximum dose: 380 mg/day

Heart Failure : Initial: 11.88 to 23.75 mg once daily; up-titrate gradually (eg, doubling the dose every 2 or more weeks) to the maximum tolerated dose while monitoring for signs and symptoms of HF; maximum dose: 190 mg/day

Hypertension : Initial : 23.75 to 95 mg once daily; titrate at weekly (or longer) intervals as needed based on patient response; maximum dose: 380 mg/day; usual dosage range: 47.5 to 190 mg once daily

Off-Label Indications:

- Secondary Prevention of Myocardial Infarction
- Maintenance of ventricular rate control in Atrial Fibrillation/ Flutter
- Maintenance therapy of Supraventricular tachycardia
- Thyrotoxicosis
- Prevention of ventricular arrhythmias
- Migraine prophylaxis

Administration:

Tablets should be taken once daily in the morning. The tablets must be swallowed whole or divided. They must not be chewed or crushed. The tablets should be taken with water.



Contraindications:

- Known hypersensitivity to product components.
- Severe bradycardia.
- Heart block greater than first degree.
- Cardiogenic shock.
- Decompensated cardiac failure.
- Sick sinus syndrome without a pacemaker.

Warnings:

Use with caution in patients with the following conditions:

Heart Failure, Bronchospastic Disease, Pheochromocytoma, Major Surgery, Diabetes, Hepatic Impairment, Thyrotoxicosis, Anaphylactic Reactions, Peripheral Vascular Disease, Myasthenia gravis, Prinzmetal variant angina.

References:

<https://www.uptodate.com>

<https://www.drugs.com>

<https://www.accessdata.fda.gov/>

<https://www.medscape.com>

<https://www.pdr.net/>